

CAMPOFORT® Special Cu

EC fertilizer

Liquid N (Mg,S) fertilizer 15 (2,5-2) with cuprum, Cu

Special guidance on application: Used in the form of outside the root nutrition by spraying on the leaves of plants. Use only when absolutely necessary, do not exceed recommended dosage.

Packaging: 24kg • 600kg • 1200kg (ie. 20l • 500l • 1000l)

Storage requirements: Store in original packaging at a temperature of -5°C to +30°C in a cool and well-ventilated place.

Maximum amount of layers during storage: 20l canister: 3 layers • IBC container: 2 layers

Shelf life: 18 months from the date of production of fertilizer under respective storage requirements

Scope and method of use: hectare dose: 12kg/ha (ie. 10l/ha)

Recommended application:

Crop	Term of application	Dosage	Possibility of combination	Water amount
cereals	BBCH 22 – 61	10l/ha (12kg/ha)	pesticides and liquid fertilizers	120 – 300l/ha
maize	BBCH 13 – 51	10l/ha (12kg/ha)	pesticides and liquid fertilizers	150 – 300l/ha
rape, mustard, poppy, sunflower	BBCH 16 – 61	10l/ha (12kg/ha)	pesticides and liquid fertilizers	120 – 300l/ha
potatoes, sugar beet	BBCH 14 – 85	10l/ha (12kg/ha)	pesticides and liquid fertilizers	150 – 300l/ha
legumes	BBCH 13 – 59	10l/ha (12kg/ha)	pesticides and liquid fertilizers	150 – 300l/ha
hops, fruits, vegetables, grapevine	according to needs	10l/ha (12kg/ha)	pesticides and liquid fertilizers	300 – 2000l/ha

Miscibility:

It is recommended to always conduct miscibility test in combination with other fertilizer and products for plant protection. Fertilizer is added as a last component. Never create tank mix from concentrated components. Add another component after perfect mixture of previous components.

Product is placed on the market in accordance with the requirements of Regulation (EC) No 2003/2003 on fertilizers of 13.10.2003 as amended. The product corresponds to the type mentioned in paragraph C.1.1. of Annex 1 to the Regulation.

Created by: AGRA GROUP a.s.

Date: 20.10.2020