

FORTE beta[®] EC fertilizer

Liquid N (Mg) fertilizer 18,4 (4,2) with boron B, manganese Mn, molybdenum Mo and zinc Zn

Special guidance on application: Used in the form of outside the root nutrition by spraying on the leaves of plants. Use only when absolutely necessary, do not exceed recommended dosage.

Packaging: 27kg • 536kg • 1340kg (ie. 20l • 400l • 1000l)

Storage requirements: Store in original packaging at a temperature of -15°C to +30°C under roof. Keep out of reach of children, away from food.

Maximum amount of layers during storage: 20l canister: 3 layers • IBC container: 2 layers

Shelf life: 18 months from the date of production of fertilizer under respective storage requirements.

Scope and method of use: hectare dose: 5,4kg/ha (ie. 4l/ha)

Recommended application:

Crop	Term of application	Dosage	Water amount
oilseeds	from 2 nd leaf until end of flowering	4l/ha (5,4kg/ha)	120 – 400l/ha
legumes	from 2 nd leaf until end of flowering	4l/ha (5,4kg/ha)	120 – 400l/ha
root crops	from 2 nd leaf until the closure of the crop	4l/ha (5,4kg/ha)	120 – 400l/ha
maize	from 2 nd leaf	4l/ha (5,4kg/ha)	120 – 400l/ha
grapevine	from 3 rd developed leaf until softening of grapes	4l/ha (5,4kg/ha)	150 – 800l/ha
fruits	from the beginning of leaf development	4l/ha (5,4kg/ha)	300 – 800l/ha
vegetables	Development of leaves or fruits	4l/ha (5,4kg/ha)	150 – 600l/ha
Hops	From 2 nd pair of leaves	4l/ha (5,4kg/ha)	1500-2000l/ha

Miscibility:

It is recommended to always conduct miscibility test in combination with other fertilizer and products for plant protection. Fertilizer is added as a last component. Never create tank mix from concentrated components.

Product is placed on the market in accordance with the requirements of Regulation (EC) No 2003/2003 on fertilizers of 13.10.2003 as amended. The product corresponds to the type mentioned in paragraph C.1.1. of Annex 1 to the Regulation.

Created by: AGRA GROUP a.s.

Date: 22.10.2020