

K-gel 175

Foliar fertilizer in form of gel

Special guidance on application: Used in the form of outside the root nutrition by spraying on the leaves of plants. Do not exceed recommended dosage.

Packaging: 25kg • 625kg • 1250kg (ie. 20l • 500l • 1000l)

Storage requirements: Store in original packaging under roof at a temperature of +5°C to +25°C in a cool and well-ventilated place. Keep out of reach of children, away from food.

Maximum amount of layers during storage: 20l canister: 3 layers • IBC container: 2 layers

Shelf life: 12 months from the date of production of fertilizer under respective storage requirements

Scope and method of use: Hectare dose might differ under existence of air bubbles in gel, therefore it can change by up to 10 %.

Recommended application:

Crop	Term of application	Number of application	Dosage	Water amount	Recommended concentration
cereals	from beginning of tillering until end of flowering	1 – 2	3 – 5 l/ha	150 – 300l/ha	1,0 – 4,0 %
maize	from 4 th leaf stage	1 – 2	3 – 5 l/ha	150 – 300l/ha	1,0 – 4,0 %
rape	from beginning of elongation growth until yellow bud phase	1 – 2	3 – 5 l/ha	150 – 300l/ha	1,0 – 4,0 %
potatoes	from 4 th leaf stage until end of August	1 – 4	3 – 5 l/ha	150 – 600l/ha	1,0 – 4,0 %
sugar beet	from 4 th leaf stage until mid-September	1 – 4	3 – 5 l/ha	150 – 600l/ha	1,0 – 4,0 %
legumes	from 5 th leaf stage until beginning of flowering	1 – 2	3 – 5 l/ha	150 – 300l/ha	1,0 – 4,0 %
grapevine	from fully developed 6 leaves until 2 weeks before harvest	1 – 4	3 – 5 l/ha	300 – 1000l/ha	0,5 – 1,0 %
hops	from height of 1m until 2 weeks before harvest	1 – 4	3 – 5 l/ha	300 – 1000l/ha	0,5 – 1,0 %
fruits	from past blossom until 2 weeks before harvest	1 – 4	3 – 5 l/ha	300 – 1000l/ha	0,5 – 1,0 %
vegetables	from beginning of intensive growth until 2 weeks before harvest	1 – 4	3 – 5 l/ha	300 – 1000l/ha	0,5 – 2,0 %
decorative plants	from beginning of intensive growth until harvest	1 – 4	3 – 5 l/ha	300 – 1000l/ha	0,5 – 2,0 %

Created by: AGRA GROUP a.s.

Date: 22.10.2020